

# Dissipationless transport in low density bilayer systems

Ady Stern<sup>a</sup>, S. Das Sarma<sup>b</sup>, Matthew P. A. Fisher<sup>c</sup> and S. M. Girvin<sup>d</sup>

(a) *Department of Condensed Matter Physics, Weizmann Institute, Rehovot 76100, Israel*

(b) *Department of Physics, University of Maryland, College Park, Maryland 20742-4111*

(c) *Institute for Theoretical Physics, UCSB, Santa Barbara, California 93106-4030*

(d) *Department of Physics, Indiana University, Bloomington, Indiana 47405-7105*

(February 1, 2008)

In a bilayer electronic system the layer index may be viewed as the  $z$ -component of an isospin- $\frac{1}{2}$ . An XY isospin-ordered ferromagnetic phase was observed in quantum Hall systems and is predicted to exist at zero magnetic field at low density. This phase is a superfluid for opposite currents in the two layers. At  $B = 0$  the system is gapless but superfluidity is not destroyed by weak disorder. In the quantum Hall case, weak disorder generates a random gauge field which probably does not destroy superfluidity. Experimental signatures include Coulomb drag and collective mode measurements.

In quantum well structures containing two separate two-dimensional electron gases in close proximity, an electron is described in terms of its position in the plane, its spin and its layer index. The latter can be regarded as an isospin- $\frac{1}{2}$ , denoted by  $\mathbf{m}$ , with the two layers being the two eigenstates of  $m_z$ . States with spontaneous XY isospin-ferromagnetic order have been observed in quantum Hall systems [1] at total Landau level filling factor  $\nu = 1$  and were predicted to exist at  $B = 0$  for sufficiently low electron density. [2] The origin of isospin ferromagnetism is a favorable Coulomb exchange energy just as in the ordinary Stoner instability.

For layer separation  $d = 0$  the isospin polarized phase breaks an SU(2) symmetry, and the problem maps onto the Stoner instability. [1] For  $d > 0$ , and in the absence of tunneling between the layers, there is an easy plane anisotropy since the direct Coulomb energy favors polarization in the XY plane ( $\langle m_z \rangle = 0$ ) in order to avoid the cost of charge imbalance between the layers that occurs for  $\langle m_z \rangle \neq 0$ . The angle of the magnetization  $\mathbf{m}(\mathbf{r})$  relative to the  $x$ -axis is then described by a field  $\varphi(\mathbf{r})$ . Because the ‘charge’ conjugate to the phase  $\varphi$  is  $m_z$ , the Goldstone mode [3] associated with the broken U(1) symmetry at finite  $d$  corresponds to superfluid currents which are *opposite* in each layer. [4,1,5]

In this paper we study transport properties of the easy plane isospin ferromagnet, focusing on the effects of disorder. At  $B = 0$  we find that disorder weakens, but does not destroy the ‘gapless isospin’ superfluidity. The lack of time-reversal symmetry in the QHE case causes disorder to induce a random gauge field which frustrates the system, but the Kosterlitz-Thouless transition probably survives weak disorder. The effect of random interlayer tunneling is a separate and different question. [6]

For  $B = 0$  it is difficult to quantify the range of parameters, particularly  $r_s$ , in which the isospin ferromagnetic state is the prevailing phase. It should lie between the low  $r_s$  paramagnetic range and the very high  $r_s$  range,

where the system forms a bilayer Wigner crystal. For a single layer, Monte Carlo calculations [7] find a ferromagnetic transition in the  $r_s \sim 20 - 30$  range, and Wigner crystallization at  $r_s \approx 37 \pm 5$  [8]. The energy differences among the various possible phases are, however, very small [7] so that a definitive statement is not possible – in fact, earlier calculations [8] did not find a ferromagnetic transition. For a double layer system, Hartree-Fock (HF) theory predicts the existence of a broken symmetry isospin ferromagnetic phase. [2] HF tends to overestimate the stability of broken symmetry states, but its predictions are often qualitatively correct and such states frequently do occur at values of  $r_s$  larger than predicted. While quantum Monte Carlo calculations are needed to obtain the precise density at which a  $B = 0$  bilayer system will undergo the spontaneous isospin ferromagnetic transition, it is reasonable to assume, based on existing HF analysis [2], that such a transition should occur at  $r_s \approx 20 - 30$ , a regime now realizable in hole systems. [9]

An HF analysis of the isospin polarized phase starts with an Hubbard-Stratanovich decomposition of the Coulomb interaction, leading to the action

$$\begin{aligned}
 S = \int dt d\mathbf{r} \Big\{ & \psi^\dagger i \partial_t \psi + \frac{1}{2m} \psi^\dagger [i \nabla - \mathbf{A}_a S^z]^2 \psi(\mathbf{r}) \\
 & - \rho_s(\mathbf{r}) V_H^{(1)}(\mathbf{r}) - \rho^z(\mathbf{r}) V_H^{(2)}(\mathbf{r}) \\
 & - V_{ex} \mathbf{m} \cdot \psi_\sigma^\dagger(\mathbf{r}) \mathbf{S}_{\sigma\sigma'} \psi_{\sigma'}(\mathbf{r}) \\
 & + \frac{1}{2} V_{ex} \mathbf{m}^2 + \frac{1}{2} \int d\mathbf{r}' n(\mathbf{r}) V_s(\mathbf{r} - \mathbf{r}') n(\mathbf{r}') \\
 & + \frac{1}{2} \int d\mathbf{r}' m_z(\mathbf{r}) V_a(\mathbf{r} - \mathbf{r}') m_z(\mathbf{r}') \Big\} \quad (1)
 \end{aligned}$$

In Eq. (1)  $\psi^\dagger, \psi$  are fermionic fields for the electrons. The symmetric and antisymmetric densities are  $\rho_s \equiv \psi_\sigma^\dagger \psi_\sigma$  and  $\rho^z \equiv \psi_\sigma^\dagger S_{\sigma\sigma'}^z \psi_{\sigma'}$ , with  $S^i$  being the Pauli matrix for the  $i$ th-component of the isospin ( $i = x, y, z$ ). The fields  $n(\mathbf{r}), \mathbf{m}(\mathbf{r})$  are auxiliary Hubbard-Stratanovich fields describing symmetric and antisymmetric densities. We are

interested in the response of the system to a weak anti-symmetric vector potential  $\mathbf{A}_a$ , which is thus included in the action (a factor of  $\frac{e}{c}$  is absorbed in  $\mathbf{A}_a$ ). In momentum representation,  $V_s(\mathbf{q}) = \frac{2\pi e^2}{q}$  and  $V_a(\mathbf{q}) = 2\pi e^2 d$  (for small  $\mathbf{q}$ ). For simplicity, we assume here that the true electron spin is fully aligned due to the Stoner instability and can be ignored (see however [10]).

In momentum representation, for small  $\mathbf{q}$ , the symmetric Hartree potential is  $V_H^{(1)} = V_s(\mathbf{q})n(\mathbf{q})$  while the antisymmetric is  $V_H^{(2)} = V_a(\mathbf{q})m_z(\mathbf{q})$ . The Fock potential  $V_{ex}$  is approximated in Eq. (1) to be local, thereby neglecting the exchange contribution to the gradient terms which contribute to the isospin stiffness. We comment on the actual value of  $V_{ex}$  and on consequences of its non-zero range below. In the system's response to  $\mathbf{A}_a$  the symmetric field  $n(\mathbf{r})$  does not play any role and we omit it from following expressions.

For fixed values of  $\mathbf{m}$  and  $n$ , the action (1) describes non-interacting electrons under the influence of a space and time dependent scalar potential  $V_H^{(1)}$ , vector potential  $\mathbf{A}_a$  and Zeeman field  $V_H^{(2)}\hat{z} + V_{ex}\mathbf{m}$ . In an  $x-y$  ordered state, the saddle point for the bosonic fields is  $n(\mathbf{r}) = m_z(\mathbf{r}) = 0$ , and  $|\mathbf{m}(\mathbf{r})| = \mathcal{M}$ , a non-zero constant. Conventional approximation schemes (HF, random phase approximation (RPA)) do not reliably obtain  $\mathcal{M}$ . Here we first assume full polarization ( $\mathcal{M} = n$ , as predicted by HF), and later discuss the case of partial polarization.

Due to an assumed lack of interlayer tunneling, the action (1) possesses a  $U(1)$  symmetry. [3] Thus, in equilibrium the system picks an arbitrary direction for  $\mathbf{m}$ . We write  $\mathbf{m} = \mathcal{M}(\cos(\varphi(\mathbf{r}))\hat{\mathbf{x}} + \sin(\varphi(\mathbf{r}))\hat{\mathbf{y}}) + m_z\hat{\mathbf{z}}$ , where  $\varphi$  is the angle between the planar component of the magnetization and the  $\hat{x}$ -axis, and expect  $\varphi$  to be constant in the ground state and slowly varying in low energy excitations. The energy cost of a deviation from the equilibrium magnetization is then expressed in terms of  $\varphi$  and  $m_z$ , and should vanish for a uniform shift in  $\varphi(\mathbf{r})$ .

We now integrate over the fermionic fields and expand the familiar  $\text{tr} \log\{\}$  term to second order in  $\varphi$  and  $m_z$ . Within RPA the expansion is given in terms of the response functions  $\chi_i \equiv -\langle \rho^i \rho^i \rangle$  and  $\chi_o \equiv -\langle \rho^z \rho^y \rangle$ . The effect of a slowly varying  $\mathbf{A}_a$  on the  $\chi$ 's can be separated out by means of a Gorkov approximation, where  $\mathbf{A}_a$  is approximated not to vary in the range of  $\mathbf{r} - \mathbf{r}'$  and  $t - t'$  in which the response function is appreciable [11]. The effect of  $\mathbf{A}_a$  is then incorporated by the “minimal coupling” prescription  $i\nabla\varphi \rightarrow (i\nabla\varphi - \mathbf{A}_a)$ , and the response functions are calculated for  $\mathbf{A}_a = 0$ . The RPA action is then (omitting the zeroth order term),

$$S_{RPA} \approx \frac{1}{2} \int d\omega \int d\mathbf{r} \left\{ \rho_s |i\nabla\varphi - \mathbf{A}_a|^2 + \frac{e^2}{\Gamma} m_z^2 + 2\chi_o V_{ex} \mathcal{M} (V_{ex} + 2\pi e^2 d) m_z \varphi \right\} \quad (2)$$

where

$$\rho_s \equiv - \lim_{\mathbf{q}, \omega \rightarrow 0} q^{-2} (1 + \chi_y V_{ex}) V_{ex} \mathcal{M}^2 \quad (3)$$

$$\frac{1}{\Gamma} \equiv - \lim_{\mathbf{q}, \omega \rightarrow 0} [1 + \chi_z (2\pi e^2 d + V_{ex})] (2\pi e^2 d + V_{ex}) \quad (4)$$

The response functions  $\chi_o, \chi_z, \chi_y$  are response functions of non-interacting electrons in a Zeeman field  $V_{ex} \mathcal{M} \hat{x}$ . For small  $\mathbf{q}, \omega$ ,

$$\begin{aligned} \chi_z = \chi_y &= -\frac{1}{2} \left[ \frac{\mathcal{M}}{\Delta - \omega - \mathcal{D}q^2} + \frac{\mathcal{M}}{\Delta + \omega - \mathcal{D}q^2} \right] \\ \chi_o &= -\frac{1}{2} \left[ \frac{i\mathcal{M}}{\Delta - \omega - \mathcal{D}q^2} - \frac{i\mathcal{M}}{\Delta + \omega - \mathcal{D}q^2} \right] \end{aligned} \quad (5)$$

where  $\Delta \equiv \mathcal{M}V_{ex}$  is the energy cost for flipping a spin, and the value of  $\mathcal{D}$  is discussed below. The  $U(1)$  invariance of the problem is the reason for  $1 + \chi_y V_{ex}$  being  $\mathcal{O}(q^2)$ . The equation of motion for  $m_z$ , derived from (2), is the Josephson type relation  $\dot{\varphi} = 2\pi e^2 d m_z$ .

The integral over  $m_z$  can now be carried out, resulting in an action in terms of  $\varphi$  and  $\mathbf{A}$  only, which is more transparent in space and time representation:

$$\int dt \int d\mathbf{r} \frac{\dot{\varphi}^2}{2} \left[ \frac{1}{2\pi e^2 d} + \frac{1}{V_{ex}} \right] + \frac{\mathcal{M}\mathcal{D}}{2} |(i\nabla - \mathbf{A}_a)\varphi|^2 \quad (6)$$

Eq. (6) is the action of a two-dimensional superfluid, with  $\mathcal{M}\mathcal{D}$  being the superfluid “spin stiffness”. *If  $\mathcal{M}\mathcal{D} > 0$ , the bilayer system responds to the vector potential  $\mathbf{A}_a$  as a superfluid. and an antisymmetric current flows without dissipation.* Eq. (6) reveals the existence of a longitudinal Goldstone mode that carries antisymmetric density and satisfies the dispersion relation

$$\omega^2 = \mathcal{M}\mathcal{D} \left[ \frac{1}{2\pi e^2 d} + \frac{1}{V_{ex}} \right]^{-1} q^2. \quad (7)$$

Within RPA, the response functions  $\chi_y, \chi_z$  are

$$\begin{aligned} \chi_y(\mathbf{q}, \omega) = \chi_z(\mathbf{q}, \omega) &= \sum_{\alpha\beta} \frac{1}{2} |\langle \alpha | \rho_{\mathbf{q}} | \beta \rangle|^2 \\ &\left\{ \frac{f(\epsilon_\alpha + \Delta) - f(\epsilon_\beta)}{\omega + \Delta + \epsilon_\alpha - \epsilon_\beta + i\eta} + \frac{f(\epsilon_\alpha) - f(\epsilon_\beta + \Delta)}{\omega - \Delta + \epsilon_\alpha - \epsilon_\beta + i\eta} \right\} \end{aligned} \quad (8)$$

where  $|\alpha\rangle, |\beta\rangle$  are single particle eigenstates of the spin-independent non-interacting Hamiltonian,  $\epsilon_\alpha, \epsilon_\beta$  are the corresponding single particle energies,  $\rho_{\mathbf{q}}$  is the density operator and  $f(\epsilon)$  is the Fermi function.

Setting  $\omega = 0$  and expanding to second order in  $q$ , we find that for a clean system with full isospin polarization ( $2\Delta > \mu$ ,  $\mu$  being the chemical potential):

$$\mathcal{M}\mathcal{D} = \int_{k < k_F} d\mathbf{k} \left[ \frac{1}{m} - \frac{k^2}{2m^2\Delta} \right] = \frac{n}{m} \left( 1 - \frac{\mu}{2\Delta} \right) \quad (9)$$

This energy cost is the sum of single particle energies of eigenstates  $|\mathbf{k}\rangle$  of electrons in a Zeeman magnetic field that precesses in space in a constant rate  $\nabla\varphi$  and is

composed of one part  $\frac{(\nabla\varphi)^2}{2m}$  originating from the anti-symmetric current induced by the precession of the field, and a second part,  $-\frac{(\mathbf{k}\cdot\nabla\varphi)^2}{2m^2\Delta}$ , which reflects the slowing down of the symmetric motion due to the field precession. There is no galilean invariance for antisymmetric currents so  $\mathcal{MD} \neq n/m$ .

The disorder potential can be separated into symmetric and antisymmetric parts. The symmetric part affects  $\mathcal{D}$  much like non-magnetic disorder does in a conventional superconductor. For weak symmetric disorder ( $k_F l \gg 1$ , and hence  $\Delta\tau \gg 1$ ), the disorder-averaged matrix elements in (8) are,

$$|\langle\alpha|\rho_{\mathbf{q}}|\beta\rangle|^2 = \frac{1}{\nu(\bar{\epsilon})} \frac{D(\bar{\epsilon})q^2}{(D(\bar{\epsilon})q^2)^2 + (\epsilon_\alpha - \epsilon_\beta)^2} \quad (10)$$

where  $\bar{\epsilon} = \frac{1}{2}(\epsilon_\alpha + \epsilon_\beta)$ ,  $D$  is the diffusion constant,  $\nu$  is the density of states, and  $|\epsilon_\alpha - \epsilon_\beta| < \frac{1}{\tau}$ . Substituting in (8) and paying attention to the dependences of  $\nu$  and  $D$  on  $\bar{\epsilon}$  we find that the effect of symmetric disorder on the spin stiffness (9) is of order  $1/\Delta\tau$ .

Antisymmetric disorder modifies the capacitive energy term in (1) to be  $2\pi e^2 d \int d\mathbf{r} [m_z(\mathbf{r}) - m_{z,dis}(\mathbf{r})]^2$  with random  $m_{z,dis}$ . As is known from studies of, e.g., Josephson junction arrays, such a randomization in the equilibrium distribution of  $m_z$  reduces the superfluid density and can, if strong enough, induce vortex-antivortex pairs destroying the superfluidity even at zero temperature. Here, since there are gapless Fermi surface excitations even in the superfluid, the resultant disordered phase may possibly be a normal Fermi liquid with no long-range inter-layer phase coherence.

Realistically, the disorder potential is made of comparable symmetric and antisymmetric components. For weak disorder, then, antisymmetric currents flow without dissipation, although the superfluid density is suppressed. As the disorder gets strong, it eventually destroys the superfluidity.

The superfluid spin stiffness  $\mathcal{MD}$  is also suppressed by finite temperature. Just as in an ordinary superconductor, its temperature dependence originates both from the Fermi functions in (8) and from thermal fluctuations of vortex-antivortex pairs in  $\varphi(\mathbf{r})$ . The spin stiffness, and with it long range order and antisymmetric dissipationless transport, disappear entirely above a Kosterlitz-Thouless (KT) transition temperature, whose precise value depends on both effects.

An experimental probe of superfluidity of antisymmetric currents is the transresistance, or drag resistance, denoted by  $\rho_D$ . In a drag measurement a current  $I_1$  is driven in one of the layers, while no current is allowed to pass through the second layer ( $I_2 = 0$ ) which develops a voltage  $V_2$ . Then,  $\rho_D \equiv -V_2/I_1$ . For two identical layers,  $\rho_D$  is the difference between the symmetric and antisymmetric resistances. In our case the latter vanishes. Thus, *the transresistance equals the symmetric one, and*

*the voltages on the two layers should be equal in magnitude and direction.* Since the superfluidity disappears at the K-T transition temperature,  $\rho_D$  would go *down* with increasing temperature. Note that for weakly coupled Fermi liquid bilayer systems  $\rho_D$  is opposite in sign to the intra-layer resistance, and its magnitude *increases* with temperature. If the superfluid mode is lost due to disorder, the antisymmetric resistance becomes appreciable, and the sign of  $\rho_D$  presumably becomes opposite to that of the intra-layer resistance.

The excitation of the sound mode (7) is another experimental probe. In the absence of isospin ferromagnetism, a double layer system has an antisymmetric acoustic plasmon mode, which is overdamped by disorder as  $q \rightarrow 0$ . [12] Here, however, the sound mode (7) is an underdamped Goldstone mode. A density sweep experiment through the transition will therefore exhibit a sharp mode at low density which will get overdamped (at long wavelengths) above the transition density. Another distinction between these two collective modes is their behavior when  $d \rightarrow 0$ . In that limit the Goldstone mode will have a long wavelength quadratic  $q^2$  dispersion, since the  $U(1)$  isospin symmetry changes into an  $SU(2)$ , whereas the normal acoustic plasmon mode tends toward the single particle dispersion  $v_F q$ .

So far we have taken the exchange Fock potential to be local, and employed RPA. The  $U(1)$  symmetry of the approximate actions (2) and (6) is exact. However, other features of our analysis are not, of which we expect two to be most important. First, as a consequence of its finite range, the Fock potential renormalizes the dispersion relation of the electrons  $\epsilon(k)$ . The sum (8) should then be evaluated with the renormalized energy dispersion, leading to the replacements  $\frac{1}{m} \rightarrow \frac{\partial^2 \epsilon}{\partial k^2}$  and  $\frac{k}{m} \rightarrow \frac{\partial \epsilon}{\partial k}$  in the integral in (9), and affecting the spin stiffness.

Second, it is conceivable that the Stoner phase is only partially isospin polarized, in contrast to the HF prediction of full isospin polarization. Interestingly, for a partially polarized state, and in the absence of disorder and electron-electron interaction, the  $q^2$ -term in the sum (8) vanishes (i.e.,  $\mathcal{D} = 0$ ), due the constant density of states  $\nu$  characteristic of two dimensions. Spin stiffness is then induced by the deviation of  $\nu$  from a constant, caused by the renormalization of the energy dispersion by interaction. Similarly, in the presence of symmetric disorder,  $\mathcal{D} \propto [(D\nu)'(\mu+\Delta/2) - (D\nu)'(\mu-\Delta/2)]$ , where a prime denotes differentiation with respect to energy. Again, the energy dispersion must deviate from parabolic for  $\mathcal{D}$  to be non-zero.

The physics of the isospin ferromagnet at filling factor  $\nu = 1$  in the QHE regime is quite different from that at  $B = 0$ . For two uncorrelated layers each at filling factor  $\nu = 1/2$ , there will be gapless ‘composite fermion’ excitations. In the presence of interlayer phase coherence however, the finite isospin stiffness leads to an energy gap

for symmetric excitations and a QHE plateau. [4,1,5]

Because of the energy gap for symmetric excitations, the fermions can be reliably integrated out [4,1,5] to yield an Euclidean action which is a functional of the isospin unit vector  $\hat{\sigma}(\mathbf{r}) \equiv \mathbf{m}(\mathbf{r})/n(\mathbf{r})$ . The Hamiltonian density corresponding to that action is

$$H = \frac{1}{2}\bar{\rho}\partial_\mu\sigma^\nu\partial_\mu\sigma^\nu + \frac{e^2}{2\Gamma}[n(\mathbf{r})\sigma^z]^2 + V_s(\mathbf{r})\delta n(\mathbf{r}) + V_a(\mathbf{r})[n_0 + \delta n(\mathbf{r})]\sigma^z(\mathbf{r}). \quad (11)$$

$\Gamma$  is the double layer capacitance per unit area (including Hartree and exchange contributions),  $V_{s,a}$  are the symmetric and antisymmetric parts of the disorder potential and  $\bar{\rho} \sim 1\text{K}$  is the exchange-induced spin stiffness. [4,5]

A crucial consequence of the quantized Hall conductivity is a constraint relating the symmetric fermion density  $n(\mathbf{r}) = n_0 + \delta n(\mathbf{r})$  to the topological (Pontryagin) density of the isospin field  $\sigma$ : [4,1,5]

$$\delta n(\mathbf{r}) = \left( \frac{\hbar}{e^2} \sigma_{xy} \right) \frac{1}{8\pi} \epsilon^{\mu\nu} \epsilon_{abc} \sigma^a \partial_\mu \sigma^b \partial_\nu \sigma^c, \quad (12)$$

where Roman indices range over  $x, y, z$  and Greek range over  $x, y$ , and  $n_0$  is the average density.

Taking advantage of the easy-plane anisotropy and noting that the XY phase angle field  $\varphi$  contains vortex singularities, we integrate out the massive  $\sigma^z$  fluctuations and find that the lack of time-reversal symmetry causes the disorder potential to generate a gauge field yielding, in the high temperature classical limit, a *2D XY model with random Dzyaloshinskii-Moriya interaction*

$$\int d\mathbf{r} \frac{\bar{\rho}}{2} |\nabla\varphi + \mathbf{a}|^2 + \sum_j [\lambda V_a(\mathbf{R}_j) - V_s(\mathbf{R}_j) Q_j] M_j \quad (13)$$

where  $Q_j = \pm 1$  is the vorticity of the  $j$ th vortex ('meron' [4,5]),  $M_j = \pm 1$  is a flavor index indicating the sign of  $\sigma^z$  in the vortex core,  $\lambda$  is a non-universal constant related to the core size, and we have dropped various irrelevant terms (e.g., a random contribution to  $\bar{\rho}$ ). The gauge field is  $\mathbf{a} \equiv \frac{\Gamma V_a}{4\pi\bar{\rho}e^2} \mathbf{J}_+$ , where  $\mathbf{J}_+ = -\frac{\hbar}{e^2} \sigma_{xy} \epsilon^{\mu\nu} \partial_\nu V_s$  is proportional to the symmetric Hall current. The  $\Gamma V_a$  term is the local density imbalance. If there is a Hall current flowing when there is a density imbalance, then more current is flowing in one layer than the other and the superfluid mode [4,5]  $\mathbf{J}_- \sim \bar{\rho}(\nabla\varphi + \mathbf{a})$  is necessarily excited. This is the physical interpretation of the gauge potential  $\mathbf{a}$  which causes these currents to flow. The phase field  $\varphi$  in (13) contains both the singular and smooth parts, and thus the first term in (13) mediates a logarithmic interaction between the merons. Unlike the  $SU(2)$  symmetric case, where a symmetric disorder potential introduces a random gauge field [13], in our case both symmetric and anti-symmetric parts are needed.

The gauge field  $\mathbf{a}$  is random with a finite correlation length. Ensemble averaging over a closed contour  $\partial\gamma$  of

perimeter  $L$  gives  $\langle \oint_{\partial\gamma} \mathbf{a} \cdot d\mathbf{r} \rangle = 0$ , and  $\langle [\oint_{\partial\gamma} \mathbf{a} \cdot d\mathbf{r}]^2 \rangle \sim L^\theta$  with  $\theta = 1$ . This is the gauge glass model for which it is known that the KT transition is destroyed in the limit of strong disorder. [14] For weak disorder the phase diagram has proven difficult to determine, [15] but it is likely that the KT transition survives. This result can be motivated by noting that in order to have an isolated flux quantum through a single plaquette, the vector potential would have to fall off like  $1/r$  and it follows that the random potential  $V_{s,a}$  would have to diverge.

To conclude, our results call for an experimental search for the  $B = 0$  phase coherent state in large  $r_s$  bilayer hole systems, and for a study of how superfluidity in this phase and its analogous phase at  $\nu = 1$  is affected by disorder. Superfluidity could be detected either in light scattering from the Goldstone mode or by drag transport.

This work was supported by grants NSF DMR-97-04005 and DMR-95-28578 (MPAF), DMR-9714055 (SMG), US-ONR (SDS), US-Israel BSF, Israeli Academy of Science, Victor Ehrlich chair (AS) and by NSF PHY94-07194 at the Institute for Theoretical Physics at UCSB.

- 
- [1] For an experimental review see: J. P. Eisenstein, Chap. 2 in *Perspectives in Quantum Hall Effects*, (Edited by Sankar Das Sarma and Aron Pinczuk (Wiley, New York, 1997). For a theoretical review see S. M. Girvin and A. H. MacDonald, Chap. 5 in the same book.
  - [2] Lian Zheng, M. W. Ortalano, and S. Das Sarma, Phys. Rev. B **55**, 4506 (1997); S. Das Sarma, M. W. Ortalano, and Lian Zheng, Phys. Rev. B **58**, 7453 (1998).
  - [3] X.-G. Wen and A. Zee, Phys. Rev. Lett. **69**, 1811 (1992).
  - [4] Kun Yang *et al.*, Phys. Rev. Lett. **72**, 732 (1994); K. Moon *et al.*, Phys. Rev. B **51**, 5138 (1995).
  - [5] S. M. Girvin, Les Houches Lectures (Springer-Verlag, Berlin and Les Editions de Physique, Paris, 1999), to be published (preprint: cond-mat/9907002).
  - [6] N. Read, Phys. Rev. B **52**, 1926 (1995).
  - [7] F. Rapisarda and G. Senatore, Aust. J. Phys. **49**, 161 (1996); D. M. Ceperley, unpublished.
  - [8] B. Tanatar and D. M. Ceperley, Phys. Rev. B **39**, 5005 (1989).
  - [9] J. Yoon *et al.*, Phys. Rev. Lett. **82**, 1744 (1999).
  - [10] S. Das Sarma, Subir Sachdev and Lian Zheng, Phys. Rev. Lett. **79**, 917 (1997); Phys. Rev. B **58**, 4672 (1998).
  - [11] P.G. deGennes, *Superfluidity of metals and alloys*, WA Benjamin (New York) 1966, chapter 7.
  - [12] S. Das Sarma and E. H. Hwang, Phys. Rev. Lett. **81**, 4216 (1998), and references therein.
  - [13] A.G. Green, Phys. Rev. Lett. **82**, 5104 (1999).
  - [14] R.A. Hyman *et al.*, Phys. Rev. B **51**, 15304 (1995).
  - [15] T. Nattermann, S. Scheidl, S. E. Korshunov, Mai Suan Li, J. de Physique I **5**, 565 (1995); C. Mudry and X.-G. Wen, Nucl. Phys. B **549**, 613 (1999).